ASPI | Afarand Scholarly Publishing Institute; Turkey
ISSN: 2626-3041 GMJ Medicine 2023;2(2):73-76 10.58209/gmjm.2.2.73

Effect of Aerobic and Anaerobic Fitness on Different Psychological Parameters in Children with Tic Disorder

ARTICLE INFO

Article Type
Original Research

Authors
Noori P.[1*]

How to cite this article
Noori P. Effect of Aerobic and Anaerobic Fitness on Different Psychological Parameters in Children with Tic Disorder. GMJ Medicine. 2023;2(2):73-76.

[1]Department of Physical Education, Kermanshah Branch, Islamic Azad University, Kermanshah, Iran

***Correspondence**
Address: -
Phone: -
Fax: -
noorip148@gmail.com

Article History
Received: October 20, 2022
Accepted: March 17, 2023
ePublished: May 3, 2023

ABSTRACT

Aims Tics are known to negatively affect psychosocial performance in children and their families. Exercise is an appropriate strategy for the prevention and management of chronic diseases. This study evaluated the effect of aerobic and anaerobic fitness on psychological function in children with tic disorder and evaluated their effects by questionnaires completed by their parents.

Materials & Methods A randomized controlled trial was carried out for 12 weeks in children with motor tics. Children were randomly assigned into three groups: aerobic, anaerobic, and Control (without exercise). The Yale global tic severity scale was used to investigate the tic severity. Parents completed Spence Children's Anxiety Scale Child and Parent Versions in the pre-test (pre-interventions) and post-test (after 12-week exercises). This study assessed the effects of aerobic and anaerobic fitness, the effect of time (pre-test & post-test), and their interaction on psychological function.

Findings The aerobic and anaerobic fitness significantly decreased Separation Anxiety, Panic and Agoraphobia, Physical Injury Fears, Social Phobia, Obsessive-Compulsive and Generalized Anxiety ($p<0.05$) compared to the control group.

Conclusion Aerobic and anaerobic fitness could alleviate psychological disorders in children with tic syndrome. Parents of children with tic can be suggested to force their children to perform aerobic and anaerobic fitness to alleviate disorders.

Keywords Aerobic Fitness; Anaerobic Fitness; Physical Injury Fears; Social Phobia; Obsessive-Compulsive; Tic Syndrome

CITATION LINKS

[1] Effect of diet, exercise and sleep on tic severity: A scoping review protocol [2] Behavioral and psychological therapies for Tourette syndrome and tic disorders: Their place in treatment and history [3] Lifetime prevalence, age of risk, and genetic relationships of comorbid psychiatric disorders in Tourette syndrome [4] Tic attacks: panic disorder in Tourette syndrome? [5] Effects of exercise and physical activity on anxiety [6] Cardiorespiratory fitness and laboratory stress: A meta-egression analysis [7] Trained men show lower cortisol, heart rate and psychological responses to psychosocial stress compared with untrained men [8] Stress and the brain: From adaptation to disease [9] Hyper-reactive hypothalamo-pituitary- adrenocortical axis in rats bred for high anxiety- related behaviour [10] 'Aerobic' and 'Anaerobic' terms used in exercise physiology: A critical terminology reflection [11] Aerobic vs anaerobic exercise training effects on the cardiovascular system [12] Indicators for high physical strain and overload in elite football players [13] Effects of aerobic and nonaerobic exercise on depression and self-concept [14] The effect of open versus closed exercises on psychological function in children with tic disorder [15] Influence of the aerobic and anaerobic training on the body fat mass in obese adolescents [16] The acute antipanic and anxiolytic activity of aerobic exercise in patients with panic disorder and healthy control subjects [17] The use of the Panic and Agoraphobia Scale (P and A) in a controlled clinical trial [18] Early exercise promotes positive hippocampal plasticity and improves spatial memory in the adult life of rats [19] The effects of an afterschool physical activity program on working memory in preadolescent children [20] Synergistic effects of diet and exercise on hippocampal function in chronically stressed mice [21] Affective beneficence of vigorous physical activity [22] Response of oxidative stress and inflammatory biomarkers to a 12-week aerobic exercise training in women with metabolic syndrome [23] Neuromodulation of aerobic exercise-a review [24] Does a 20-week aerobic exercise training programme our capabilities to buffer real-life stressors? A randomized, controlled trial using ambulatory assessment [25] DHEA, DHEA-S and cortisol responses to acute exercise in older adults in relation to exercise training status and sex [26] Salivary cortisol and testosterone responses to resistance and plyometric exercise in 12- to 14-year-old boys [27] The impacts of physical exercise on stress coping and well-being in university students in the context of leisure

Copyright© 2023, the Authors | Publishing Rights, ASPI. This open-access article is published under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License which permits Share (copy and redistribute the material in any medium or format) and Adapt (remix, transform, and build upon the material) under the Attribution-NonCommercial terms.

Introduction

Tourette syndrome (TS) is known as a neurodevelopmental disorder in patients with motor and vocal tics for at least one year [1]. Tics are not serious disorders but transient disorders in childhood that influence children's development [2]. Tics are short and insignificant, such as blinking, grimacing head, nose twitching, mouth, and shoulder movements [1]. Tics are short, but these influence psychological performance in children and members of the family [2]. Children with tic disorders are faced with neuropsychiatric conditions, such as attention deficit hyperactivity disorder and obsessive-compulsive [3]. It was reported that 85% of anxiety disorders in children with tic disorders [3]. Parents and children commonly report distressed behaviors associated with faulted daily function and anxiety disorders [4]. Different strategies are used for decreasing psychological symptoms in patients with tic syndrome, such as drugs, physical activity, etc. The application of drugs has been faced with limitations due to their side effects [2]. Researchers are interested in using physical activity to alleviate psychological disorders. It was reported that using exercises alleviates diseases [5].

Physical activity is an appropriate strategy for the alleviation of adverse effects of disorders. Exercise is considered the first strategy for the prevention and management of disorders. Performing regular exercise causes physiological changes and adaptations in the human body by affecting hypothalamic-pituitary-adrenal (HPA) axis reactivity [6, 7]. The region has an important role in improving adaptive responses to physical and psychological stressors [8]. It was reported depressive and anxiety signs in patients with faults in the HPA [9]. Exercises are divided into aerobic and anaerobic exercises [10]. Aerobic and anaerobic exercises have different effects on the physiology of the body. It was reported that this type of exercise can appropriately affect mental indices [11]. Other studies did not find positive effects of physical activities on mental indices [12]. It was reported that the efficiency of anaerobic exercises in decreasing anxiety [13].

Seemingly, aerobic and anaerobic fitness can have positive effects on psychological function in children with motor tics. Psychological disorders are a part of tic syndrome that are not considered by the scientific community, but these influence children. To the best of our knowledge, no study has been conducted to evaluate the effect of aerobic and anaerobic fitness on psychological function in children with a tic disorder. This study was thus conducted to evaluate the effect of aerobic and anaerobic fitness on separation anxiety, generalized anxiety, social phobia, obsessive-compulsive behaviors, panic and agoraphobia, and physical injury fears in children with a tic disorder.

Materials and Methods

Ethics Statement

All the procedures used were in agreement with the local Ethical Committee of the Physical Education Review Board of Islamic Azad University, Kermanshah-Iran (No. of permission, IAUKI-13401). We collected consent from parents of all the subjects before their children participated. Not only children but also their parents completed consent forms.

Study procedure

This randomized controlled trial was conducted for 12 weeks in children with motor tics. Forty-five girl children were randomly assigned into 3 groups: 1) the Aerobic group, where children exercised aerobic fitness for 12 weeks; 2) the Anaerobic group, where children exercised anaerobic exercises for 12 weeks; and 3) the Control group, where children did not receive any training.

Participants

Forty-five children ranging from 9 to 11 years (10.78±1.24) were studied in this study and completed evaluations at post-test. The flowchart of the study is illustrated in Figure 1. Body weight (in shorts and t-shirt) and height (without shoes) were investigated by an electronic scale and converted to body mass index (BMI). Children were randomly selected from local elementary schools in the Kermanshah Province, West region. The inclusion criteria were as follows; 1) Consent of children for participation in the study, 2) Children aged between 10 and 12 years of age, 3) Children had no medical condition that would affect study results or limit physical activity, 4) Children with tic severity >2 and 5) Children were not taking medication that would affect study results (e.g. antipsychotics). Exclusion criteria included: 1) Children did not have a tendency to participate in this study, 2) Children with tic severity <2, 3) Consumption of drugs that would affect study results or limit physical activity, 4) The use of antipsychotics, and 5) Non-participating in exercises for at least 2 meetings.

Yale Global Tic Severity Scale

The Yale global tic severity scale is a clinician-rated measure for tic severity over the last 7–10 days that has a stable factor structure and excellent psychometric properties. The motor and tics are rated at a 0–5 scale across 5 dimensions: number, frequency, intensity, complexity, and interference [14].

Exercise intervention

The exercise sessions were carried out after school and were conducted in consecutive sessions. Children were transported by bus to a training place in Kermanshah City. Instructors included master's level physical educators, bachelor's level exercise scientists, and other trained research staff who had at least some college education. In this study, we emphasized enjoyment and safety but no competition. The aerobic training program was

designed using a cycle ergometer, triplicated a week, for 12 weeks, and for an initial period of 40 minutes [16]. The anaerobic training was performed using the method of "Interspaced Training [15]. In this study, parents completed Spence Children's Anxiety Scale Child and Parent Versions in the pre-test (pre-interventions) and post-test (after 12-week exercises). The questionnaire included 38 items rated on a 4-point scale (0 to 3; never to always), and the SCAS-C includes six additional positive filler items. The SCAS-C/P comprises six subscales addressing Separation Anxiety (six items), Generalized Anxiety (six items), Social Phobia (six items), Obsessive–Compulsive Behaviors (six items), Panic and Agoraphobia (nine items), and Physical Injury Fears (five items).

Statistical Analysis

The present study was conducted in a 2×3 factorial arrangement with the effect of time (pre-test & post-test) and exercises (Control, Aerobic & Anaerobic) and their interaction. The data were reported as mean±SD and analyzed for the main effects of time and exercise and the interactions between time and exercise. The data were analyzed using SPSS 23 software. Comparisons between groups were conducted by the Sceffe test, and the T-test conducted paired-tests. Kolmogorov-Smirnov test was used to test the normality.



Figure 1. Flowchart of the study

Findings

Descriptive characteristics

There were no significant differences between groups (p>0.05) according to age, weight, height, and BMI conditions.

Table 1. The subjects' characteristics

Parameter	Control	Aerobic	Anaerobic	p-Value
Age (Year)	10.56±0.85	11.23±0.85	10.57±0.83	0.905
Weight (Kg)	33.12±1.35	32.21±1.91	34.51±2.12	0.823
Height (m)	1.29±0.10	1.32±0.21	1.27±0.11	0.821
BMI (Kg/m^2)	19.95±2.26	18.51±2.20	21.43±2.11	0.356

Yale Global Tic Severity Scale

Table 2 presents our findings for five dimensions in the different groups. The results did not show significant differences between groups (p>0.05).

Table 2. Distribution of Yale global tic severity scale scores across participants

Parameter	Control	Aerobic	Anaerobic	p-Value
Number	3.32±0.41	3.57±0.71	3.61±0.42	0.783
Frequency	4.21±0.57	4.10±0.57	4.32±0.59	0.571
Intensity	3.12±0.51	3.09±0.52	3.01±0.58	0.711
Complexity	2.62±0.72	2.57±0.62	2.42±0.51	0.712
Interference	2.57±0.49	2.51±0.43	2.39±0.41	0.523

Normality

Table 3 shows the normality distribution results for all the parameters in the pre-test and post-test. The results showed that values higher than 0.05 meant normality.

Table 3. The results for normality status (all were normal)

Parameter	Pre-test		Post-test	
	Z	P-value	Z	P-value
Separation Anxiety	1.282	0.081	1.083	0.079
Generalized Anxiety	1.115	0.212	0.911	0.314
Social Phobia	1.241	0.059	0.859	0.341
Obsessive–Compulsive	1.029	0.239	1.002	0.091
Panic and Agoraphobia	1.222	0.152	1.226	0.071
Physical Injury Fears	0.751	0.326	1.112	0.099

The effect of aerobic and anaerobic fitness on Separation Anxiety

The effect of aerobic and anaerobic fitness on Separation Anxiety for time ($F_{(1, 44)}$=7.10; p=0.0009), fitness ($F_{(2, 44)}$=5.41; p=0.001) and time × fitness interaction ($F_{(1, 44)}$=6.12; p=0.001) were shown in Table 4. Scheffe test showed a significant difference between groups in the post-test (p=0.0001). The results showed that aerobic and anaerobic fitness decreased separation anxiety in the post-test (Table 4).

Table 4. The effect of aerobic and anaerobic fitness on Separation Anxiety in pre and post-Test

Groups	Pre-test	Post-test	T-test
Control	2.61±0.59	2.65±0.44	0.731
Aerobic	2.64±0.48	2.11±0.47	0.029
Anaerobic	2.65±0.39	2.08±0.48	0.036
Scheffe	0.912	0.0001	

The effect of aerobic and anaerobic fitness on Panic and Agoraphobia

The effect of aerobic and anaerobic fitness on Panic and Agoraphobia for time ($F_{(1, 44)}$=7.08; p=0.0001), fitness ($F_{(2, 44)}$=7.22; p=0.0001) and time × fitness interaction ($F_{(1, 44)}$=7.67; p=0.0001) were shown in Table 5. There was a significant response for the Scheffe test in the post-test (p=0.0001). The results showed that aerobic and anaerobic fitness reduced Panic and Agoraphobia (Table 5).

Table 5. Effect of Fitness on Panic and Agoraphobia in pre and post-Test

Groups	Pre-test	Post-test	T-test
Control	2.85±0.53	2.83±0.69	0.912
Aerobic	2.91±0.72	2.13±0.21	0.0001
Anaerobic	2.79±0.50	2.05±0.24	0.0001
Scheffe	0.812	0.0001	

The effect of aerobic and anaerobic fitness on Physical Injury Fears

The effect of aerobic and anaerobic fitness on Physical Injury Fears for time ($F_{(1, 44)}$=31.48; p=0.0001), fitness ($F_{(2, 44)}$=15.48; p=0.0001) and time × fitness interaction ($F_{(1, 44)}$=16.39; p=0.0001) were shown in Table 6. Scheffe test showed a significant difference between groups in the post-test (p=0.0001). Aerobic and anaerobic fitness decreased Physical Injury Fears (Table 6).

Table 6. Effect of skill exercises on Physical Injury Fears in pre and post-Test

Groups	Pre-test	Post-test	T-test
Control	2.89±0.51	2.78±0.50	0.956
Aerobic	2.75±0.51	1.32±0.44	0.0001
Anaerobic	2.74±0.31	1.42±0.44	0.0001
Scheffe	0.923	0.00001	

The effect of aerobic and anaerobic fitness on Generalized Anxiety

The effect of aerobic and anaerobic fitness on Generalized Anxiety for time ($F_{(1, 44)}$=14.20; p=0.0001), fitness ($F_{(2, 44)}$=21.18; p=0.0001), and time × fitness interaction ($F_{(1, 44)}$=15.49; p=0.0001) were shown in Table 7. There was a significant response for the Scheffe test in the post-test (p=0.001).

Table 7. The effects of skill exercises on Generalized Anxiety in pre and post-Test

Groups	Pre-test	Post-test	T-test
Control	2.83±0.21	2.75±0.51	0.759
Aerobic	2.79±0.34	2.05±0.21	0.001
Anaerobic	2.71±0.35	2.08±0.22	0.001
Scheffe	0.938	0.001	

The effect of aerobic and anaerobic fitness on Social Phobia

The effect of aerobic and anaerobic fitness on Social Phobia for time ($F_{(1, 44)}$=19.60; p=0.0001), fitness ($F_{(2, 44)}$=21.45; p=0.0001), and time × fitness interaction ($F_{(1, 44)}$=23.12; p=0.0001) were shown in Table 8. Significant responses were observed for the Scheffe test in the post-test (p=0.0001).

Table 8. The effects of skill exercises on aerobic and anaerobic fitness on Social Phobia in pre and post-Test

Groups	Pre-test	Post-test	T-test
Control	2.91±0.12	2.75±0.51	0.759
Aerobic	2.82±0.25	2.05±0.21	0.001
Anaerobic	2.85±0.25	2.08±0.22	0.001
Scheffe	0.901	0.0001	

The effect of aerobic and anaerobic fitness on Obsessive–Compulsive

The effect of aerobic and anaerobic fitness on Social Phobia for time ($F_{(1, 44)}$=21.20; p=0.0001), fitness ($F_{(2, 44)}$=23.31; p=0.0001), and time × fitness interaction ($F_{(1, 44)}$=19.75; p=0.0001) were shown in Table 9. Significant responses were observed for the Scheffe test in the post-test (p=0.0001).

Table 9. The effects of skill exercises on aerobic and anaerobic fitness on Obsessive–Compulsive in pre and post-Test

Groups	Pre-test	Post-test	T-test
Control	2.86±0.22	2.91±0.22	0.759
Aerobic	2.87±0.32	2.02±0.14	0.001
Anaerobic	2.91±0.19	2.01±0.09	0.001
Scheffe	0.901	0.0001	

Discussion

The results showed that aerobic and anaerobic fitness had positive effects on decreasing psychological disorders in children with tic syndrome. No study has been found to show the effects of aerobic and anaerobic fitness on psychological parameters in children with motor tics. Conversely, Strohle *et al.* [16] assessed the effects of aerobic treadmill exercise on anti-panic and anxiolytic activity in patients with panic disorder and showed that anxiety was significantly higher in patients submitted to treadmill exercise. The results

are in contrast with our findings. The reason for the difference between our findings and others could be attributed to physiological conditions. Parallel to our findings, Bandelow *et al.* [17] aerobic exercise decreased anxiety symptoms. In another study, Gomes da Silva [18] showed that physical activity affects attention, depression, anxiety, and overall well-being. Other studies showed that aerobic and anaerobic fitness improved neuropsychological and electrophysiological performances of working memory in preadolescent children [19]. An animal study assessed the effects of long-term wheel running on anxiety-like behavior in transgenic mice and showed decreased anxiety-like behavior [20]. It was reported that aerobic exercises have positive effects on the density of neural conveyors in the neural system and reduce muscular tension and hormone levels [20-25]. Other studies showed that aerobic exercises decrease the level of stress and anxiety by influencing the automatic neural system and decreasing anxiety and depression [25]. Aerobic fitness decreases cortisol levels and anxiety [26]. Kim *et al.* [27] investigated the responses of five participants to aerobic exercises and anaerobic exercises and showed that different fitness improved mental health and happiness, each with a different degree, which verifies the results of the present study. Other studies showed that aerobic and group activities are more effective for improving mental well-being compared to anaerobic and individual exercises [25].

Conclusion

Both fitness similarly decreased psychological disorders in children with tic syndrome. It can be suggested to use aerobic and aerobic fitness to decrease psychological disorders in patients with psychological disorders.

Acknowledgments: This work was supported by Islamic Azad University, Kermanshah Branch, Kermanshah, Iran.
Ethical Permission: None declared by the authors.
Conflicts of Interests: None declared by the authors.
Funding/Support: None declared by the authors.

References

1- Nosratmirshekarlou E, Shafiq S, Goodarzi Z.S, Martino D, Pringsheim T. Effect of diet, exercise and sleep on tic severity: A scoping review protocol. BMJ Open. 2019;9:e024653.
2- Robertson MM, Chowdhury U. Behavioral and psychological therapies for Tourette syndrome and tic disorders: Their place in treatment and history. Neuropsychiatry. 2011;1(4): 385-400.
3- Hirschtritt ME, Lee PC, Pauls DL, Dion Y, Grados MA, Cornelia Illman, *et al*. Lifetime prevalence, age of risk, and genetic relationships of comorbid psychiatric disorders in Tourette syndrome. JAMA Psychiatry. 2015;72:325-33.
4- Robinson S, Hindley P, Hedderly T. Tic attacks: panic disorder in Tourette syndrome? Poster on 19th International Congress of Parkinson's Disease and Movement Disorders; 2015 June 14-18; San Diego, CA. Milwaukee: MDS; 2016.
5- Anderson E, Shivakumar G. Effects of exercise and physical activity on anxiety. Frontier in Psychiatry. 2013;4:1-4.
6- Jackson EM, Dishman RK. Cardiorespiratory fitness and laboratory stress: A meta-egression analysis. Psychophysiology. 2006;43:57-72.
7- Rimmele U, Zellweger BC, Marti B, Seiler R, Mohiyeddini C, Ehlert U, et al. Trained men show lower cortisol, heart rate, and psychological responses to psychosocial stress compared with untrained men. Psychoneuroendocrinology. 2007;32:627-35.
8- De Kloet ER, Joëls M, Holsboer F. Stress and the brain: From adaptation to disease. Natural Review Neuroscience. 2005;6:463-75.
9- Landgraf R, Wigger A, Holsboer F, Neumann I. Hyper-reactive hypothalamic-pituitary-adrenocortical axis in rats bred for high anxiety-related behavior. J Neuroendocrinol. 1999;11(6):405-7.
10- Chamari K, Padulo J. 'Aerobic' and 'Anaerobic' terms used in exercise physiology: A critical terminology reflection. Sports Med Open. 2015;1(1):9.
11- Patel H, Alkhawam H, Madanieh R, Shah N, Kosmas CE, Vittorio TJ. Aerobic vs anaerobic exercise training effects on the cardiovascular system. World J Cardiol. 2017;9(2):134-8.
12- Meister S, Faude O, Ammann T, Schnittker R, Meyer T. Indicators for high physical strain and overload in elite football players. Scand J Med Sci Sports. 2013;23(2):156-63.
13- Stein PN, Motta RW. Effects of aerobic and nonaerobic exercise on depression and self-concept. Percept Mot Skills. 1992;74(1):79-89.
14- Shiravand M, Safikhani H. The effect of open versus closed exercises on psychological function in children with tic disorder. GMJ Medicine. 2020;4(1):196-205.
15- Cláudia Fernandez A, de Mello MT, Tufik S, de Castro PM, Fisberg M. Influence of the aerobic and anaerobic training on the body fat mass in obese adolescents. Rev Bras Med Esporte. 2004;10:159-65.
16- Ströhle A, Graetz B, Scheel M, Wittmann A, Feller C, Heinz A, et al. The acute anti-panic and anxiolytic activity of aerobic exercise in patients with panic disorder and healthy control subjects. J Psychiatr Res. 2009;43(12):1013-7.
17- Bandelow B, Broocks A, Pekrun G, George A, Meyer T, Pralle L, et al. The use of the Panic and Agoraphobia Scale (P and A) in a controlled clinical trial. Pharmacopsychiatry. 2000;33(5):174-81.
18- Gomes da Silva S, Unsain N, Hugo Mascó D, Toscano-Silva M, Alves de Amorim H, Henrique Silva Araújo B, et al. Early exercise promotes positive hippocampal plasticity and improves spatial memory in the adult life of rats. Hippocampus. 2012;22:347-58.
19- Kamijo K, Pontifex MB, O'Leary KC, Scudder MR, Wu CT, Castelli DM, et al. The effects of an afterschool physical activity program on working memory in preadolescent children. Dev Sci. 2011;14:1046-58.
20- Hutton CP, Déry N, Rosa E, Lemon JA, Rollo CD, Boreham DR, et al. Synergistic effects of diet and exercise on hippocampal function in chronically stressed mice. Neuroscience. 2015;308:180-93.
21- Morgan WP. Affective beneficence of vigorous physical activity. Med Sci Sports Exerc. 1985;17(1):94-100.

22- Farinha JB, Steckling FM, Stefanello ST, Cardoso MS, Nunes LS, Barcelos RP, et al. Response of oxidative stress and inflammatory biomarkers to a 12-week aerobic exercise training in women with metabolic syndrome. Sports Med Open. 2015;1(1):19.
23- Heijnen S, Hommel B, Kibele A, Colzato LS. Neuromodulation of aerobic exercise-a review. Front Psychol. 2016;6:1890.
24- von Haaren B, Ottenbacher J, Muenz J, Neumann R, Boes K, Ebner-Priemer U. Does a 20-week aerobic exercise training program our capabilities to buffer real-life stressors? A randomized, controlled trial using ambulatory assessment. Eur J Appl Physiol. 2016;116(2):383-94.
25- Heaney JL, Carroll D, Phillips AC. DHEA, DHEA-S and cortisol responses to acute exercise in older adults in relation to exercise training status and sex. Age (Dordr). 2013;35(2):395-405.
26- Klentrou P, Giannopoulou A, McKinlay BJ, Wallace P, Muir C, Falk B, et al. Salivary cortisol and testosterone responses to resistance and plyometric exercise in 12- to 14-year-old boys. Appl Physiol Nutr Metab. 2016;41(7):714-8.
27- Kim JH, McKenzie LA. The impacts of physical exercise on stress coping and well-being in university students in the context of leisure. Health. 2014;6(19):2570-80.